



TITLE:

A GEOMETRIC INTERPRETATION OF ISOPARAMETRIC HYPERSURFACES (Geometry of homogeneous spaces and submanifolds)

AUTHOR(S):

Kimura, Makoto

CITATION:

Kimura, Makoto. A GEOMETRIC INTERPRETATION OF ISOPARAMETRIC HYPERSURFACES (Geometry of homogeneous spaces and submanifolds). 数理解析研究所講究録 1998, 1044: 1-3

ISSUE DATE:

1998-04

URL:

<http://hdl.handle.net/2433/62147>

RIGHT:

A GEOMETRIC INTERPRETATION OF ISOPARAMETRIC HYPERSURFACES

木村真琴 (MAKOTO KIMURA)

Department of Mathematics
Faculty of Education
Ibaraki University

1. Isoparametric hypersurfaces in a real space form.

This part is a joint work with Sadahiro Maeda [KM]. In differential geometry it is interesting to know the shape of a Riemannian submanifold by observing the extrinsic shape of geodesics of the submanifold. For example: A hypersurface M^n isometrically immersed into a real space form $\widetilde{M}^{n+1}(c)$ of constant curvature c (that is, $\widetilde{M}^{n+1}(c) = \mathbb{R}^{n+1}$, $S^{n+1}(c)$ or $H^{n+1}(c)$ according as the curvature c is zero, positive, or negative) is totally umbilic in $\widetilde{M}^{n+1}(c)$ if and only if every geodesic of M , considered as a curve in the ambient space $\widetilde{M}^{n+1}(c)$, is a circle. Here we treat a geodesic as a circle of null curvature.

In this talk we are interested in a hypersurface M^n of a real space form $\widetilde{M}^{n+1}(c)$ satisfying that there exists such an *orthonormal* basis $\{v_1, \dots, v_n\}$ at each point p of the hypersurface M^n that all geodesics of M^n through p in the direction v_i ($1 \leq i \leq n$) are circles in the ambient space $\widetilde{M}^{n+1}(c)$. The classification problem of such hypersurfaces is concerned with studies about isoparametric hypersurfaces M^n 's in a real space form $\widetilde{M}^{n+1}(c)$ (that is, all principal curvatures of M^n in $\widetilde{M}^{n+1}(c)$ are constant).

Theory of isoparametric submanifolds is one of the most interesting objects in differential geometry. In particular, É. Cartan studied extensively isoparametric hypersurfaces in a standard sphere. The classification problem of isoparametric hypersurfaces in a sphere is still open (for details, see [CR]).

The initial purpose of this talk is to provide a characterization of all isoparametric hypersurfaces by observing the extrinsic shape of geodesics of hypersurfaces in a real space form.

Theorem 1. *Let M^n be a connected hypersurface of a real space form $\widetilde{M}^{n+1}(c)$ of constant curvature c . Then M^n is isoparametric in $\widetilde{M}^{n+1}(c)$ if and only if there exists such an orthonormal basis $\{v_1, \dots, v_m\}$ of the orthogonal complement of $\ker A$ in $T_p(M)$ ($m = \text{rank } A$) that all geodesics of M through p in the direction v_i ($1 \leq i \leq m$) are circles of nonzero curvature in the ambient space $\widetilde{M}^{n+1}(c)$.*

*This research was partially supported by Grant-in-Aid for Scientific Research (No. 09740050), Ministry of Education, Science and Culture.

Theorem 2. *Let M^n be a connected hypersurface of a real space form $\widetilde{M}^{n+1}(c)$ of constant curvature c . Then M^n is isoparametric with nonzero constant principal curvatures in $\widetilde{M}^{n+1}(c)$ if and only if for each point p of M , there exists such an orthonormal basis $\{v_1, \dots, v_n\}$ of $T_p(M)$ that all geodesics of M through p in the direction v_i ($1 \leq i \leq n$) are circles of nonzero curvature in the ambient space $\widetilde{M}^{n+1}(c)$.*

Theorem 3. *Let M^n be a connected hypersurface of a real space form $\widetilde{M}^{n+1}(c)$ of constant curvature c . Then M^n is isoparametric in $\widetilde{M}^{n+1}(c)$ if and only if for each point p of M , there exists such an orthonormal basis $\{v_1, \dots, v_n\}$ of $T_p(M)$ of principal curvature vectors that all geodesics of M through p in the direction v_i ($1 \leq i \leq n$) are circles in the ambient space $\widetilde{M}^{n+1}(c)$.*

2. Homogeneous real hypersurfaces in a complex projective space.

This part is a joint work with Toshiaki Adachi and Sadahiro Maeda [AKM]. Let $P_n(\mathbb{C})$ be an n -dimensional complex projective space with Fubini-Study metric of constant holomorphic sectional curvature 4, and let M be a real hypersurface of $P_n(\mathbb{C})$. Then M has an almost contact metric structure (ϕ, ξ, η, g) inherited from the Kaehler structure of $P_n(\mathbb{C})$. Many differential geometers have studied M by using this structure (cf. [O]). Typical examples of real hypersurfaces in $P_n(\mathbb{C})$ are homogeneous real hypersurfaces, that is, real hypersurfaces given as orbits under subgroups of the projective unitary group $PU(n+1)$.

Takagi ([T]) classified homogeneous real hypersurfaces in $P_n(\mathbb{C})$. Due to his work, we find that a homogeneous real hypersurface in $P_n(\mathbb{C})$ is locally congruent to one of the six model spaces of type A_1, A_2, B, C, D and E . They are realized as tubes of constant radius over compact Hermitian symmetric spaces of rank 1 or 2. A homogeneous real hypersurface of type A_1 is usually called a *geodesic hypersphere*.

In the study of real hypersurfaces in $P_n(\mathbb{C})$, there can be the following two problems:

- (A) Give a characterization of homogeneous real hypersurfaces in $P_n(\mathbb{C})$.
- (B) Construct non-homogeneous nice real hypersurfaces in $P_n(\mathbb{C})$ and characterize such examples.

In this talk we are interested in Problem (A). In differential geometry it is interesting to know the shape of a Riemannian submanifold by observing the extrinsic shape of geodesics of the submanifold. From this point of view we here recall the fact that a hypersurface M^n in \mathbb{R}^{n+1} is locally a standard sphere if and only if all geodesics of M are circles of positive curvature in \mathbb{R}^{n+1} . We shall provide a characterization of all homogeneous real hypersurfaces in $P_n(\mathbb{C})$ by observing the shape of geodesics on the real hypersurfaces as curves in $P_n(\mathbb{C})$.

The purpose of this part is to prove the following result which is an improvement of the previous paper [MO].

Theorem 4. *Let M be a connected real hypersurface of $P_n(\mathbb{C})$. Then M is congruent to a homogeneous real hypersurface if and only if there exist such orthonormal vectors $v_1, v_2, \dots, v_{2n-2}$ orthogonal to ξ at each point p of M that all geodesics $\gamma_i = \gamma_i(s)$ on M with $\gamma_i(0) = p$ and $\dot{\gamma}_i(0) = v_i$ ($1 \leq i \leq 2n-2$) are circles in $P_n(\mathbb{C})$ with positive curvature.*

In the hypothesis of our Theorem we do not need to suppose that we take the vectors $\{v_1, \dots, v_{2n-2}\}$ as a local field of orthonormal frames in M . However, for all homogeneous real hypersurfaces M' s in $P_n(\mathbb{C})$, we can take a local field of orthonormal frames in M satisfying the hypothesis of our Theorem.

It is well-known that there does not exist a real hypersurface all of whose geodesics are circles in $P_n(\mathbb{C})$. Every circle in Theorem is a simple closed curve which lies on some totally real totally geodesic $P^2(\mathbb{R})$ in $P_n(\mathbb{C})$. We note that for any homogeneous real hypersurface M , at each point p of M the geodesic $\gamma = \gamma(s)$ with $\gamma(0) = p$ and $\dot{\gamma}(0) = \xi$ is also a circle in $P_n(\mathbb{C})$ which is a simple closed curve lying on some holomorphic totally geodesic $P_1(\mathbb{C})$ in $P_n(\mathbb{C})$. All circles in $P_n(\mathbb{C})$ are simple curves. However, a circle in $P_n(\mathbb{C})$ is not necessarily closed (see, [AMU]).

REFERENCES

- [AKM] T. Adachi, M. Kimura and S. Maeda, *A characterization of all homogeneous real hypersurfaces in a complex projective space by observing the extrinsic shape of geodesics*, preprint.
- [AMU] T. Adachi, S. Maeda and S. Udagawa, *Circles in a complex projective space*, Osaka J. Math. **32** (1995), 709–719.
- [CR] T. E. Cecil and P. J. Ryan, *Tight and Taut immersions of manifolds*, Research Notes in Mathematics, vol. 107, Pitman, Boston, 1985.
- [K] M. Kimura, *Real hypersurfaces and complex submanifolds in complex projective space*, Trans. Amer. Math. Soc. **296** (1986), 137–149.
- [KM] M. Kimura and S. Maeda, *Geometric meaning of isoparametric hypersurfaces in a real space form*, preprint.
- [MO] S. Maeda and K. Ogiue, *Characterizations of geodesic hyperspheres in a complex projective space by observing the extrinsic shape of geodesics*, Math. Z. **225** (1997), 537–542.
- [O] M. Okumura, *On some real hypersurfaces and complex submanifolds in a complex projective space*, Trans. Amer. Math. Soc. **212** (1975), 355–364.
- [T] R. Takagi, *On homogeneous real hypersurfaces of a complex projective space*, Osaka J. Math. **10** (1973), 495–506.

BUNKYO, MITO, IBARAKI, 310, JAPAN

E-mail address: mkimura@mito.ipc.ibaraki.ac.jp